

A Weaker Constraint Qualification of Globally Convergent Homotopy Method for a Multiobjective Programming Problem

Guangming Yao¹, Wen Song²

¹Department of Mathematics, Clarkson University, Potsdam, USA

²Department of Mathematics, Harbin Normal University, Harbin, China
 Email: gyao@clarkson.edu, wsong218@yahoo.com.cn

Received September 29, 2012; revised January 9, 2013; accepted January 16, 2013

ABSTRACT

In this paper, we prove that the combined homotopy interior point method for a multiobjective programming problem introduced in Ref. [1] remains valid under a weaker constrained qualification—the Mangasarian-Fromovitz constrained qualification, instead of linear independence constraint qualification. The algorithm generated by this method associated to the Karush-Kuhn-Tucker points of the multiobjective programming problem is proved to be globally convergent.

Keywords: Multiobjective Programming Problem; Homotopy Method; KKT Condition; Efficient Solution; MFCQ

1. Introduction

Let \mathbb{R}^n be the n -dimensional Euclidean space, and let \mathbb{R}_+^n and \mathbb{R}_{++}^n denote the nonnegative and positive \mathbb{R}^n , respectively. For any two vectors $y = (y_1, y_2, \dots, y_n)$ and $z = (z_1, z_2, \dots, z_n)$ in \mathbb{R}^n , we use the following conventions: $y = z$ iff $y_i = z_i, i = 1, 2, \dots, n$. Similarly, we can define $y \leq z, y < z$, and $y \leq z$.

Consider the following multiobjective programming problem (MOP)

$$\begin{aligned} \min f(x) \\ \text{s.t. } g(x) \leq 0, h(x) = 0, \end{aligned}$$

where

$$\begin{aligned} x \in \mathbb{R}^n, f = (f_1, f_2, \dots, f_p)^T, \\ g = (g_1, g_2, \dots, g_m)^T, h = (h_1, h_2, \dots, h_s)^T. \end{aligned}$$

We assume that all $f_i, i = 1, \dots, p, g_j, j \in I$ and $h_k, k \in J$ are twice continuously differentiable functions, where $I = \{1, 2, \dots, m\}, J = \{1, 2, \dots, s\}$.

Let

$$\begin{aligned} \Omega &= \{x \in \mathbb{R}^n : g(x) < 0, h(x) = 0\}, \\ \Omega_1 &= \{x \in \mathbb{R}^n : g(x) < 0\}, \\ \Lambda^{++} &= \left\{ \lambda \in \mathbb{R}_+^p : \sum_{i=1}^p \lambda_i = 1 \right\}, \\ I(x) &:= \{j \in I : g_j(x) = 0\}. \end{aligned}$$

It is well known that if x is an efficient solution of

(MOP), under some constraint qualifications, such as the Kuhn and Tucker constraint qualification (see Ref. [2]) or the Abadie constraint qualification (see Ref. [3]), then the following Karush-Kuhn-Tucker (KKT) condition at x for (MOP) holds (see Refs. [4,5]):

$$\begin{cases} \nabla f(x)^T \lambda + \nabla g(x)^T u + \nabla h(x)^T v = 0 \\ Ug(x) = 0 \\ h(x) = 0 \end{cases} \quad (1)$$

where $\lambda \in \mathbb{R}_+^p \setminus \{0\}, u \in \mathbb{R}_+^m, v \in \mathbb{R}^s$, and

$$U = \text{diag}\{u_1, u_2, \dots, u_m\}.$$

We say that x is a KKT point of (MOP) if it satisfies the KKT condition.

Since the remarkable papers of Kellogg *et al.* (Ref. [6]) and Chow *et al.* (Ref. [7]) have been published, more and more attention has been paid to the homotopy method. As a globally convergent method, the homotopy method (or path-following method) now becomes an important tool for numerically solving nonlinear problems including nonlinear mathematical programming and complementarity problems (see Refs. [3,4]).

In 1988, Megiddo (see Ref. [8]) and Kojima *et al.* (see Ref. [9]) discovered that the Karmakar interior point method was a kind of path-following method for solving linear programming. Since then, the interior path-following method has been generalized to convex programming, and becomes one of the main methods for solving mathematical programming problems. Among most interior

methods, one of the main ideas is numerically tracing the center path generated by the optimal solution set of the so-called logarithmic barrier function. Usually, the strict convexity of the logarithmic barrier function or non-emptiness and boundedness of the feasible set (see Ref. [10]) are needed. In 1997, Lin, Yu and Feng (see Ref. [11]) presented a new interior point method—combined homotopy interior point method (CHIP method)—for convex nonlinear programming without such assumptions. Subsequently, Lin, Li and Yu (see Ref. [12]) generalized CHIP method to general nonlinear programming where, instead of convexity condition, they used a

more general “normal cone condition”.

In 2003, Lin, Zhu and Sheng (see Ref. [13]) generalized CHIP method to convex multiobjective programming (CMOP) with only inequality constraints. Instead of (CMOP), they considered an associated non-convex nonlinear scalar optimization problem and constructed the homotopy mapping.

In Refs. [1,14], we considered a combined homotopy interior point method for the multiobjective programming (MOP) under the condition linearly independent constraint qualification (LICQ). To find a KKT point of (MOP), we construct a homotopy as follows

$$0 = H(\omega, \omega^0, \mu) = \begin{bmatrix} (1-\mu)(\nabla f(x)^T \lambda + \nabla g(x)^T u) + \nabla h(x)^T v + \mu(x - x^0) \\ h(x) \\ Ug(x) - \mu U^0 g(x^0) \\ (1-\mu)\left(1 - \sum_{i=1}^p \lambda_i\right)e - \mu(\lambda^{2/5} - (\lambda^0)^{2/5}) \end{bmatrix} \tag{2}$$

where $\omega^0 = (x^0, \lambda^0, u^0, v^0) \in \Omega \times \Lambda^{++} \times R_{++}^m \times \{0\}$, $e = (1, 1, \dots, 1)^T \in R^p$, $\omega = (x, \lambda, u, v) \in \bar{\Omega} \times R_{++}^{p+m} \times R^s$, $\mu \in [0, 1]$, $\lambda^{2/5} = (\lambda_1^{2/5}, \dots, \lambda_p^{2/5})$, and $(\lambda^0)^{2/5} = ((\lambda_1^0)^{2/5}, \dots, (\lambda_p^0)^{2/5})$.

Let

$$H_{\omega^0}^{-1}(0) = \{(\omega, \mu) \in \Omega \times R_{++}^{p+m} \times R^s \times (0, 1] : H(\omega, \omega^0, \mu) = 0\}.$$

Let $A \subset R^n$ be a nonempty closed set and $x \in A$. We recall that the Fréchet normal cone of A at x is defined as

$$N_A(x) = \left\{ x^* \in R^n \mid \limsup_{x' \xrightarrow{A} x} \frac{\langle x^*, x' - x \rangle}{\|x' - x\|} \leq 0 \right\}.$$

We used the following basic assumptions which are commonly used in that literature:

(A1) Ω is nonempty (Slater condition) and bounded;

(A2) (LICQ) $\forall x \in \bar{\Omega}$, the matrix

$$(\nabla h(x)^T, \nabla g_j(x)^T : j \in I(x))$$

is a matrix of full column rank;

(A3) Normal condition:

$$\forall x \in \bar{\Omega}, \{x + \hat{N}_{\bar{\Omega}}(x)\} \cap \bar{\Omega} = \{x\}.$$

It is well known that if condition (A2) holds, then

$$\hat{N}_{\bar{\Omega}}(x) = \left\{ \sum_{i \in I(x)} u_i \nabla g_i(x) + \sum_{j \in J} v_j \nabla h_j(x) : u_i \geq 0, i \in I(x), v \in R^s \right\}. \tag{3}$$

We have proved the following convergence result in Ref. [1].

Theorem 1.1 (Convergence of the method) Suppose f_i, g_j , and h_k are twice continuously differentiable functions such that the conditions (A1), (A2), and (A3) hold. Then for almost all

$$\omega^0 \in \Omega \times \Lambda^{++} \times R_{++}^m \times R^s,$$

the zero-point set $H_{\omega^0}^{-1}(0)$ of the homotopy map (2) contains a smooth curve $\Gamma_{\omega^0} \subset \bar{\Omega} \times R_{++}^{p+m+s} \times (0, 1]$, which starts from $(\omega^0, 1)$. As $\mu \rightarrow 0$, the limit set

$T \times \{0\} \subset \bar{\Omega} \times R_{++}^{p+m+s} \times \{0\}$ of Γ_{ω^0} is nonempty, and the x -component of every point in T is a KKT point of (MOP).

Recently, many researchers extended and improved the results in Ref. [1] to convex multiobjective programming problem, see Ref. [14-17]. The purpose of this paper is to show that Theorem 1.1 remains true under the condition MFCQ instead of LICQ. The paper is organized as following. In Section 2, we prove the existence and convergence of a smooth homotopy path from almost any interior initial point ω^0 to a solution of

the KKT system of (MOP) under the condition MFCQ.

2. Main Results

We need the following elementary condition.

(A2') (MFCQ) For every $x \in \bar{\Omega}$, the following conditions hold:

- $\nabla h_j(x), j \in J$, are linear independent;
- there exists a $p \in R^n$ such that

$$\nabla g_i(x)p > 0, i \in I(x) \text{ and } \nabla h_j(x)p = 0, j \in J.$$

Clearly, condition (A2) implies (A2'). It is also known that if (A2') holds, then (3) remains valid.

By using an analogue argument as in Ref. [1], we can prove the following two theorems.

Theorem 2.1 Suppose that $\Omega \neq \emptyset$ and conditions (A1), (A2') hold. Then for almost all initial points $\omega^0 \in \Omega \times \Lambda^{++} \times \mathbb{R}_{++}^m \times \{0\}$, 0 is a regular value of H_{ω^0} and $H_{\omega^0}^{-1}(0)$ consists of some smooth curves. Among them, a smooth curve, say Γ_{ω^0} starts from $(\omega^0, 1)$.

Theorem 2.2 Suppose that $\Omega \neq \emptyset$ and conditions (A1), (A2') hold. For a given $\omega^0 \in \Omega \times \Lambda^{++} \times \mathbb{R}_{++}^m \times \mathbb{R}^s$, if 0 is a regular value of H_{ω^0} , then the projection of the smooth curve Γ_{ω^0} on the λ component is bounded.

We next prove that Γ_{ω^0} is a bounded curve.

Theorem 2.3 (Boundedness) Suppose that the conditions (A1), (A2'), and (A3) hold. Then for a given $\omega^0 \in \Omega \times \Lambda^{++} \times \mathbb{R}_{++}^m \times \mathbb{R}^s$, if 0 is a regular value of H_{ω^0} , then Γ_{ω^0} is a bounded curve.

Proof: By Theorem 2.2, it is sufficient to show that the (u, v) -component of smooth curve is bounded. Suppose that there exists a sequence $\{(\omega^k, \mu_k)\} \subset \Gamma_{\omega^0}$,

such that

$$x^k \rightarrow x^*, \mu_k \rightarrow \mu_*, \lambda^k \rightarrow \lambda^*$$

and

$$\|(u^k, v^k)\| \rightarrow \infty (k \rightarrow \infty),$$

where $x^* \in \Omega, \mu_* \in [0, 1], \lambda^* \geq 0$. Since closed unit circle of R^{m+s} is compact, without loss of generality we can assume that

$$\lim_{k \rightarrow \infty} \frac{(u^k, v^k)}{\|(u^k, v^k)\|} = (u^*, v^*).$$

Clearly, $u^* \geq 0, \|(u^*, v^*)\| = 1$. By (2), we have

$$(1 - \mu_k) \left(\nabla f(x^k)^T \lambda^k + \nabla g(x^k)^T u^k \right) \tag{4}$$

$$+ \nabla h(x^k)^T v^k + \mu_k (x^k - x^0) = 0,$$

$$U^k g(x^k) - \mu_k U^0 g(x^0) = 0. \tag{5}$$

Let

$$I_1(x^*) = \left\{ i \in I : \lim_{k \rightarrow \infty} u_i^k = \infty \right\},$$

$$I_2(x^*) = \left\{ j \in J : \lim_{k \rightarrow \infty} v_j^k = \infty \right\}.$$

By (5), we know $I_1(x^*) \subset I(x^*)$.

Rewrite (4) as

$$(1 - \mu_k) \left[\nabla f(x^k)^T \lambda^k + \sum_{i \in I(x^*)} u_i^k \nabla g_i(x^k) \right. \tag{6}$$

$$\left. + \sum_{i \in I(x^*)} u_i^k \nabla g_i(x^k) \right] + \nabla h(x^k)^T v^k + \mu_k (x^k - x^0) = 0.$$

Divide (6) by $\|(u^k, v^k)\|$ and let $k \rightarrow \infty$, since $\|(u^k, v^k)\| \rightarrow \infty$, (6) becomes

$$\nabla g_I(x^*)^T ((1 - \mu_*)u^*) + \nabla h(x^*)^T v^* = 0, \tag{7}$$

where

$$\nabla g_I(x^*) = \left\{ (g_i(x^*) : i \in I(x^*)) \right\},$$

$$u^* = \left\{ (u_i^* : i \in I(x^*)) \right\}.$$

1) If $u_i^* = 0$, then $u^* = 0$, and $\nabla h(x^*)^T v^* = 0$. By (A2'), $v^* = 0$. This is a contradiction with $\|(u^*, v^*)\| = 1$.

2) If $u_i^* \neq 0$, we consider the following two cases:

1. If $\mu_* \in [0, 1)$, we know $(1 - \mu_*)u_i^* \geq 0$ because of $u_i^* \neq 0$. By (A2'), there exists a nonzero vector $p \in R^n$, such that

$$p^T \nabla g_I(x^*)^T > 0 \text{ and } p^T \nabla h(x^*)^T = 0. \tag{8}$$

This, together with (7), implies that $(1 - \mu_*)u_i^* = 0$, which is a contradiction.

2. If $\mu_* = 1$, by (7) and (A2'), we know $v^* = 0$. So,

$$\|u^*\| = 1 \text{ since } \|(u^*, v^*)\| = 1.$$

Because of (5), $I_1(x^*) = I(x^*) \neq \emptyset$. Thus

$$x^* \in \partial \Omega_1, \|u_i^k\| \rightarrow \infty (k \rightarrow \infty), \forall i \in I.$$

Without loss of generality, we can assume that

$$u_i^k > 0, \mu_k \neq 1 (\forall k \in N), \text{ and } \lim_{k \rightarrow \infty} \frac{u_i^k}{\|u_i^k\|} = u.$$

Hence $u \geq 0, \|u\| = 1$.

a) If $I_2(x^*) = \emptyset$, then $\{v^k\}$ is bounded. We may assume $\lim_{k \rightarrow \infty} v^k = v$. Divide (6) by $\|(1 - \mu_k)u_i^k\|$ and let $k \rightarrow \infty$, (6) becomes

$$\begin{aligned} & \nabla g_I(x^*)^T u \\ & + \lim_{k \rightarrow \infty} \left[\frac{1}{\|(1-\mu_k)u_I^k\|} \left(\nabla h(x^k)^T v^k + \mu_k(x^k - x^0) \right) \right] = 0. \end{aligned} \tag{9}$$

This implies that

$$\lim_{k \rightarrow \infty} \frac{1}{\|(1-\mu_k)u_I^k\|}$$

exists. Indeed,

$$\begin{aligned} & \lim_{k \rightarrow \infty} \left[\nabla h(x^k)^T v^k + \mu_k(x^k - x^0) \right] \\ & = \nabla h(x^*)^T v + x^* - x^0. \end{aligned}$$

If

$$\nabla h(x^*)^T v + x^* - x^0 = 0,$$

that is

$$\nabla h(x^*)^T v + x^* = x^0.$$

By condition (A3), $x^* = x^0$. This is impossible since $x^* \in \partial\Omega$. So

$$\nabla h(x^*)^T v + x^* - x^0 \neq 0.$$

By (9), we then have that $\lim_{k \rightarrow \infty} \frac{1}{\|(1-\mu_k)u_I^k\|}$ exists.

Assume $\lim_{k \rightarrow \infty} \frac{1}{\|(1-\mu_k)u_I^k\|} = \mu.$

Then $\mu \geq 0$.

If $\mu > 0$, (9) becomes

$$\nabla g_I(x^*)^T \frac{u}{\mu} + \nabla h(x^*)^T v + x^* = x^0.$$

This contradicts to condition (A3).

If $\mu = 0$, (9) becomes

$$\nabla g_I(x^*)^T u = 0.$$

By (A2'), there exists a nonzero vector $p \in R^n$ such that

$$\nabla g_I(x^*)^T p > 0, \text{ and } p^T \nabla g_I(x^*)^T u = 0.$$

Thus $u = 0$, which contradicts $\|u\| = 1$.

b) If $I_2(x^*) \neq \emptyset$, without loss of generality, we can assume that

$$\lim_{k \rightarrow \infty} \frac{\left((1-\mu_k)u_I^k, v^k \right)}{\left\| \left((1-\mu_k)u_I^k, v^k \right) \right\|} = (\hat{u}, \hat{v}),$$

where $\|(\hat{u}, \hat{v})\| = 1$. Since

$$\left\| \left((1-\mu_k)u_I^k, v^k \right) \right\| \rightarrow \infty \text{ as } k \rightarrow \infty, \tag{10}$$

we divide (4) by $\left\| \left((1-\mu_k)u_I^k, v^k \right) \right\|$, and let $k \rightarrow \infty$, we have that

$$\nabla g_I(x^*)^T \hat{u} + \nabla h(x^*)^T \hat{v} = 0. \tag{11}$$

If $\hat{u} = 0$, then

$$\nabla h(x^*)^T \hat{v} = 0.$$

By condition (A2'), $\hat{v} = 0$. This is a contradiction since $\|(\hat{u}, \hat{v})\| = 1$.

If $\hat{u} \neq 0$, by (A2'), there is a nonzero vector $p \in R^n$ such that

$$p^T \nabla g_I(x^*)^T > 0, \text{ and } p^T \nabla h(x^*)^T = 0.$$

This, together with (11), implies $\hat{u} = 0$. This is a contradiction.

Therefore, Γ_{ω^0} is a bounded curve.

By an analogue argument as in Ref. [1], it is easy to show the following result.

Theorem 2.4 (Convergence of the method) *Suppose that the conditions (A1), (A2'), and (A3) hold. Then for almost all $\omega^0 \in \Omega \times \Lambda^{++} \times \mathbb{R}_{++}^m \times \mathbb{R}^s$, the zero-point set $H_{\omega^0}^{-1}(0)$ of the homotopy map (2) contains a smooth curve $\Gamma_{\omega^0} \subset \bar{\Omega} \times \mathbb{R}_+^{p+m} \times (0,1]$, which starts from $(\omega^0, 1)$. As $\mu \rightarrow 0$, the limit set $T \times \{0\} \subset \bar{\Omega} \times \mathbb{R}_+^{p+m} \times \{0\}$ of Γ_{ω^0} is nonempty, and every point in T is a solution of (1).*

Therefore, Theorem 2.4 shows that for almost all $\omega^0 \in \Omega \times \mathbb{R}_+^{p+m} \times \mathbb{R}^s \times \{0\}$, the homotopy Equation (2) generates a smooth curve Γ_{ω^0} starts from $(\omega^0, 1)$ which is called the homotopy path, the limit set $T \times \{0\} \subset \bar{\Omega} \times \mathbb{R}_+^{p+m+s} \times \{0\}$ of Γ_{ω^0} is nonempty, and the x -component of every point in T is a KKT point of (MOP), the ω -component of the homotopy path is the solution of (1) as μ goes to 0.

REFERENCES

- [1] W. Song and G. M. Yao, "Homotopy Method for General Multiobjective Programming Problems," *Journal of Optimization Theory and Applications*, Vol. 138, No. 1, 2008, pp. 139-153. doi:10.1007/s10957-008-9366-6
- [2] H. W. Kuhn and A. W. Tucker, "Nonlinear Programming," *Proceedings of the Second Berkeley Symposium on Mathematical Statistics and Probability*, University of California Press, Berkeley, 1951.
- [3] T. Maeda, "Second-Order Conditions for Efficiency in Nonsmooth Multiobjective Optimization Problems," *Journal of Optimization Theory and Applications*, Vol. 122, No. 3, 2004, pp. 521-538.

- [doi:10.1023/B:JOTA.0000042594.46637.b4](https://doi.org/10.1023/B:JOTA.0000042594.46637.b4)
- [4] C. Y. Lin and J. L. Dong, "Methods and Theories in Multiobjective Optimization," Jinlin Education Press, Changchun, 1992.
- [5] M. Abadie, "Generalized Kuhn-Tucker Conditions for Mathematical Programming," *SIAM Journal on Control*, Vol. 7, No. 2, 1969, pp. 232-241. [doi:10.1137/0307016](https://doi.org/10.1137/0307016)
- [6] R. B. Kellogg, T. Y. Li and J. A. Yorke, "A Constructive Proof of the Brouwer Fixed-Point Theorem and Computational Results," *SIAM Journal on Numerical Analysis*, Vol. 13, No. 4, 1976, pp. 473-483. [doi:10.1137/0713041](https://doi.org/10.1137/0713041)
- [7] S. N. Chow, J. Mallet-Paret and J. A. Yorke, "Finding Zeros of Maps: Homotopy Methods That are Constructive with Probability One," *Mathematical Computation*, Vol. 32, 1978, pp. 887-899. [doi:10.1090/S0025-5718-1978-0492046-9](https://doi.org/10.1090/S0025-5718-1978-0492046-9)
- [8] N. Megiddo, "Pathways to the Optimal Set in Linear Programming, in Progress in Mathematical Programming, Interior Point and Related Methods," Springer, New York, 1988, pp. 131-158.
- [9] M. Kojima, S. Mizuno and A. Yoshise, "A Primal-Dual Interior Point Algorithm for Linear Programming," In: N. Megiddo, Ed., *Progress in Mathematical Programming, Interior Point and Related Methods*, Springer, New York, 1988, pp. 29-47.
- [10] E. L. Allgower and K. Georg, "Numerical Continuation Methods: An Introduction," Springer Verlag, Berlin, 1990. [doi:10.1007/978-3-642-61257-2](https://doi.org/10.1007/978-3-642-61257-2)
- [11] Z. H. Lin, B. Yu and G. C. Feng, "A Combined Homotopy Interior Method for Convex Nonlinear Programming," *Applied Mathematics and Computation*, Vol. 84, No. 2-3, 1997, pp. 193-211. [doi:10.1016/S0096-3003\(96\)00086-0](https://doi.org/10.1016/S0096-3003(96)00086-0)
- [12] Z. H. Lin, Y. Li and B. Yu, "A Combined Homotopy Interior Point Method for General Nonlinear Programming Problems," *Applied Mathematics and Computation*, Vol. 80, No. 2-3, 1996, pp. 209-224. [doi:10.1016/0096-3003\(95\)00295-2](https://doi.org/10.1016/0096-3003(95)00295-2)
- [13] Z. H. Lin, D. L. Zhu and Z. P. Sheng, "Finding a Minimal Efficient Solution of a Convex Multiobjective Program," *Journal of Optimization Theory and Applications*, Vol. 118, No. 1, 2003, pp. 587-600. [doi:10.1023/A:1024739508603](https://doi.org/10.1023/A:1024739508603)
- [14] Y. F. Shang and B. Yu, "A Constraint Shifting Homotopy Method for Convex Multi-Objective Programming," *Journal of Computational and Applied Mathematics*, Vol. 236, No. 5, 2011, pp. 640-646.
- [15] X. Zhao, S. G. Zhang and Q. H. Liu, "Homotopy Interior-Point Method for a General Multiobjective Programming Problem," *Journal of Applied Mathematics*, Vol. 2012, 2012, Article ID: 497345.
- [16] N. Kim and L. Thuy, "An Algorithm for Generating Efficient Outcome Points for Convex Multiobjective Programming Problem," *Intelligent Information and Database Systems Lecture Notes in Computer Science*, Vol. 5991, No. 2010, 2010, pp. 390-399.
- [17] Z. Chen, "Multiobjective Optimization Problems, Vector Variational Inequalities and Proximal-Type Methods," Dissertations, Hong Kong Polytechnic University, Kowloon, 2010.